

For the Use of a Registered Medical Practitioner Only

1. GENERIC NAME AND BRAND NAME

Bempedoic Acid and Ezetimibe Tablets 180 mg/10 mg
Bemneo® EZ

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film coated tablet contains

Bempedoic acid.....180 mg

Ezetimibe I.P.10mg

Excipients..... q.s.

Colours: Brilliant Blue FCF Aluminium Lake, Indigo carmine Aluminium Lake and Titanium Dioxide I.P.

3. DOSAGE FORM AND STRENGTH

Refer section 1 & 2

4. CLINICAL PARTICULARS

4.1 THERAPEUTIC INDICATION

It is indicated as an adjunct to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or established atherosclerotic cardiovascular disease who required additional lowering of LDL C.

4.2 POSOLOGY AND METHOD OF ADMINISTRATION

Posology

The recommended dose of Bempedoic acid + ezetimibe is one film-coated tablet of 180 mg/10 mg taken once daily.

Coadministration with bile acid sequestrants

Dosing of Bempedoic acid + ezetimibe should occur either at least 2 hours before or at least 4 hours after administration of a bile acid sequestrant.

Concomitant simvastatin therapy

When Bempedoic acid + ezetimibe is coadministered with simvastatin, simvastatin dose should be limited to 20 mg daily (or 40 mg daily for patients with severe hypercholesterolaemia and high risk for cardiovascular complications, who have not achieved their treatment goals on lower doses and when the benefits are expected to outweigh the potential risks)

Special populations

Elderly patients: No dose adjustment is necessary in elderly patients

Patients with renal impairment: No dose adjustment is necessary in patients with mild or moderate renal impairment. There are limited data available in patients with severe renal impairment (defined as estimated glomerular filtration rate [eGFR] < 30 mL/min/1.73 m²), and patients with end-stage renal disease (ESRD) on dialysis have not been studied with bempedoic acid. Additional monitoring for adverse reactions may be warranted in these patients when Bempedoic acid + ezetimibe is administered.

Patients with hepatic impairment: No dose adjustment is necessary in patients with mild hepatic impairment (Child-Pugh A). Treatment with Bempedoic acid + ezetimibe is not recommended in patients with moderate (Child-Pugh B) or severe (Child-Pugh C) hepatic impairment due to the unknown effects of the increased exposure to ezetimibe.

Paediatric population: The safety and efficacy of Bempedoic acid + ezetimibe in children aged less than 18 years have not been established. No data are available. Method of administration Each film-coated tablet should be taken orally with or without food. Tablet should be swallowed whole

4.3 CONTRAINDICATIONS

- Hypersensitivity to the active substances or to any of the excipients.
- Pregnancy.
- Breast-feeding.
- Concomitant use with simvastatin > 40 mg daily.
- Bempedoic acid + ezetimibe coadministered with a statin is contraindicated in patients with active liver disease or unexplained persistent elevations in serum transaminases.

4.4.SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Potential risk of myopathy with concomitant use of statins

Bempedoic acid increases plasma concentrations of statins. Statins occasionally cause myopathy. In rare cases, myopathy may take the form of rhabdomyolysis with or without acute renal failure secondary to myoglobinuria, and can lead to fatality. In postmarketing experience with ezetimibe, very rare cases of myopathy and rhabdomyolysis have been reported. Most patients who developed rhabdomyolysis were taking a statin concomitantly with ezetimibe.

Patients receiving Bempedoic acid + ezetimibe as adjunctive therapy to a statin should be monitored for adverse reactions that are associated with the use of high doses of statins. All patients receiving Bempedoic acid + ezetimibe in addition to a statin should be advised of the potential increased risk of myopathy and told to report promptly any unexplained muscle pain, tenderness, or weakness. If such symptoms occur while a patient is receiving treatment with Bempedoic acid + ezetimibe and a statin, a lower maximum dose of the same statin or an alternative statin, or discontinuation of Bempedoic acid + ezetimibe and initiation of an alternative lipid-lowering therapy should be considered under close monitoring of lipid levels and adverse reactions. If myopathy is confirmed by a creatine phosphokinase (CPK) level > 10× upper limit of normal (ULN), Bempedoic acid + ezetimibe and any statin that the patient is taking concomitantly should be immediately discontinued.

Myositis with a CPK level > 10× ULN was rarely reported with bempedoic acid and background simvastatin 40 mg therapy. Doses of simvastatin > 40 mg should not be used with Bempedoic acid + ezetimibe.

Tendon Rupture

Bempedoic acid, a component of Bempedoic acid + ezetimibe, is associated with an increased risk of tendon rupture or injury. In clinical trials, tendon rupture occurred in 0.5% of patients treated with bempedoic acid versus 0% of placebo-treated patients and involved the rotator cuff (the shoulder), biceps tendon, or Achilles tendon. Tendon rupture occurred

within weeks to months of starting bempedoic acid. Tendon rupture may occur more frequently in patients over 60 years of age, in those taking corticosteroid or fluoroquinolone drugs, in patients with renal failure, and in patients with previous tendon disorders. Discontinue Bempedoic acid + ezetimibe immediately if the patient experiences rupture of a tendon. Consider discontinuing Bempedoic acid + ezetimibe if the patient experiences joint pain, swelling, or inflammation. Advise patients to rest at the first sign of tendinitis or tendon rupture and to contact their healthcare provider if tendinitis or tendon rupture symptoms occur. Consider alternative therapy in patients with a history of tendon disorders or tendon rupture.

Increased serum uric acid

Bempedoic acid may raise the serum uric acid level due to inhibition of renal tubular OAT2 and may cause or exacerbate hyperuricaemia and precipitate gout in patients with a medical history of gout or predisposed to gout. Treatment with Bempedoic acid + ezetimibe should be discontinued if hyperuricaemia accompanied with symptoms of gout appear.

Elevated liver enzymes

In clinical trials, elevations of $> 3 \times$ ULN in the liver enzymes alanine aminotransferase (ALT) and aspartate aminotransferase (AST) have been reported with bempedoic acid. These elevations have been asymptomatic and not associated with elevations $\geq 2 \times$ ULN in bilirubin or with cholestasis and have returned to baseline with continued treatment or after discontinuation of therapy. In controlled coadministration trials in patients receiving ezetimibe with a statin, consecutive transaminase elevations ($\geq 3 \times$ ULN) have been observed. Liver function tests should be performed at initiation of therapy. Treatment with Bempedoic acid + ezetimibe should be discontinued if an increase in transaminases of $> 3 \times$ ULN persists.

Renal impairment

There is limited experience with bempedoic acid in patients with severe renal impairment (defined as $\text{eGFR} < 30 \text{ mL/min/1.73 m}^2$), and patients with ESRD on dialysis have not been studied with bempedoic acid. Additional monitoring for adverse reactions may be warranted in these patients when Bempedoic acid + ezetimibe is administered.

Hepatic impairment

Due to the unknown effects of the increased exposure to ezetimibe in patients with moderate to severe hepatic impairment (Child-Pugh B and C), Bempedoic acid + ezetimibe is not recommended in these patients

Fibrates

The safety and efficacy of ezetimibe administered with fibrates have not been established. If cholelithiasis is suspected in a patient receiving Bempedoic acid + ezetimibe and fenofibrate, gallbladder investigations are indicated and this therapy should be discontinued.

Ciclosporin

Caution should be exercised when initiating Bempedoic acid + ezetimibe in the setting of ciclosporin. Ciclosporin concentrations should be monitored in patients receiving Bempedoic acid + ezetimibe and ciclosporin.

Anticoagulants

If Bempedoic acid + ezetimibe is added to warfarin, other coumarin anticoagulants, or fluindione, the International Normalised Ratio (INR) should be appropriately monitored.

Contraception

Women of childbearing potential must use effective contraception during treatment. Patients should be advised to stop taking Bempedoic acid + ezetimibe before stopping contraceptive measures if they plan to become pregnant.

4.5 DRUGS INTERACTIONS

No specific pharmacokinetic drug interaction studies with Bempedoic acid + ezetimibe have been conducted. Drug interactions that have been identified in studies with bempedoic acid or ezetimibe determine the interactions that may occur with Bempedoic acid + ezetimibe.

Effects of other medicinal products on individual components of Bempedoic acid + ezetimibe

Fibrates

Concomitant fenofibrate or gemfibrozil administration modestly increased total ezetimibe concentrations (approximately 1.5- and 1.7-fold, respectively). Fenofibrate may increase cholesterol excretion into the bile, leading to cholelithiasis. In a preclinical study in dogs, ezetimibe increased cholesterol in the gallbladder bile. A lithogenic risk associated with the therapeutic use of Bempedoic acid + ezetimibe cannot be ruled out.

If cholelithiasis is suspected in a patient receiving Bempedoic acid + ezetimibe and fenofibrate, gallbladder studies are indicated and alternative lipid-lowering therapy should be considered.

Ciclosporin

In a study of eight post-renal transplant patients with creatinine clearance of > 50 mL/min on a stable dose of ciclosporin, a single 10 mg dose of ezetimibe resulted in a 3.4-fold (range 2.3- to 7.9-fold) increase in the mean area under the curve (AUC) for total ezetimibe compared to a healthy control population, receiving ezetimibe alone, from another study (n=17). In a different study, a renal transplant patient with severe renal impairment who was receiving ciclosporin and multiple other medicinal products demonstrated a 12-fold greater exposure to total ezetimibe compared to concurrent controls receiving ezetimibe alone. In a two-period crossover study in twelve healthy subjects, daily administration of 20 mg ezetimibe for 8 days with a single 100 mg dose of ciclosporin on day 7 resulted in a mean 15% increase in ciclosporin AUC (range 10% decrease to 51% increase) compared to a single 100 mg dose of ciclosporin alone. A controlled study on the effect of coadministered ezetimibe on ciclosporin exposure in renal transplant patients has not been conducted. Caution should be exercised when initiating Bempedoic acid + ezetimibe in the setting of ciclosporin. Ciclosporin concentrations should be monitored in patients receiving Bempedoic acid + ezetimibe and ciclosporin.

Cholestyramine

Concomitant cholestyramine administration decreased the mean AUC of total ezetimibe (ezetimibe plus ezetimibe glucuronide) approximately 55%. The incremental low-density

lipoprotein cholesterol (LDL-C) reduction due to adding Bempedoic acid + ezetimibe to cholestyramine may be lessened by this interaction.

Transporter-mediated drug interactions

In vitro drug interaction studies suggest bempedoic acid, as well as its active metabolite and glucuronide form, are not substrates of commonly characterised drug transporters with the exception of bempedoic acid glucuronide, which is an OAT3 substrate.

Probenecid

Probenecid, an inhibitor of glucuronide conjugation, was studied to evaluate the potential effect of these inhibitors on the pharmacokinetics of bempedoic acid. Administration of bempedoic acid 180 mg with steady-state probenecid resulted in a 1.7-fold increase in bempedoic acid AUC and a 1.9-fold increase in bempedoic acid active metabolite (ESP15228) AUC. These elevations are not clinically meaningful and do not impact dosing recommendations.

Effects of individual components of Bempedoic acid + ezetimibe on other medicinal products

Statins

The pharmacokinetic interactions between bempedoic acid 180 mg and simvastatin 40 mg, atorvastatin 80 mg, pravastatin 80 mg, and rosuvastatin 40 mg were evaluated in clinical trials. Administration of a single dose of simvastatin 40 mg with steady-state bempedoic acid 180 mg resulted in a 2-fold increase in simvastatin acid exposure. Elevations of 1.4-fold to 1.5-fold in AUC of atorvastatin, pravastatin, and rosuvastatin (administered as single doses) and/or their major metabolites were observed when coadministered with bempedoic acid 180 mg. Higher elevations have been observed when these statins were coadministered with a supratherapeutic 240 mg dose of bempedoic acid.

No clinically significant pharmacokinetic interactions were seen when ezetimibe was coadministered with atorvastatin, simvastatin, pravastatin, lovastatin, fluvastatin or rosuvastatin.

Transporter-mediated drug interactions

Bempedoic acid and its glucuronide weakly inhibit OATP1B1 and OATP1B3 at clinically relevant concentrations. Coadministration of Bempedoic acid + ezetimibe with medicinal products that are substrates of OATP1B1 or OATP1B3 (i.e., bosentan, fimasartan, asunaprevir, glecaprevir, grazoprevir, voxilaprevir, and statins such as atorvastatin, pravastatin, fluvastatin, pitavastatin, rosuvastatin, and simvastatin may result in increased plasma concentrations of these medicinal products.

Bempedoic acid inhibits OAT2 in vitro, which may be the mechanism responsible for minor elevations in serum creatinine and uric acid. Inhibition of OAT2 by bempedoic acid may also potentially increase plasma concentrations of medicinal products that are substrates of OAT2. Bempedoic acid may also weakly inhibit OAT3 at clinically relevant concentrations.

Anticoagulants Concomitant administration of ezetimibe (10 mg once daily) had no significant effect on bioavailability of warfarin and prothrombin time in a study of twelve healthy adult males. However, there have been postmarketing reports of increased INR in patients who had ezetimibe added to warfarin or fluindione.

If Bempedoic acid + ezetimibe is added to warfarin, other coumarin anticoagulants, or fluindione, INR should be appropriately monitored.

Other interactions studied

Bempedoic acid had no effect on the pharmacokinetics of oral contraceptive norethindrone/ethinyl estradiol. In clinical interaction studies, ezetimibe had no effect on the pharmacokinetics of oral contraceptives ethinyl estradiol and levonorgestrel. Bempedoic acid had no effect on the pharmacokinetics or pharmacodynamics of metformin.

In clinical interaction studies, ezetimibe had no effect on the pharmacokinetics of dapson, dextromethorphan, digoxin, glipizide, tolbutamide, or midazolam, during coadministration.

4.6 USE IN SPECIAL POPULATIONS (SUCH AS PREGNANT WOMEN, LACTATING WOMEN, PAEDIATRIC PATIENTS, GERIATRIC PATIENTS ETC.)

Pregnancy

Bempedoic acid + ezetimibe is contraindicated during pregnancy.

There are no or limited amount of data from the use of Bempedoic acid + ezetimibe in pregnant women. Studies in animals with bempedoic acid have shown reproductive toxicity.

Because bempedoic acid decreases cholesterol synthesis and possibly the synthesis of other cholesterol derivatives needed for normal foetal development, Bempedoic acid + ezetimibe may cause foetal harm when administered to pregnant women. Bempedoic acid + ezetimibe should be discontinued prior to conception or as soon as pregnancy is recognized.

Women of childbearing potential

Women of childbearing potential should use effective contraception during treatment.

Breast-feeding

It is unknown whether bempedoic acid/metabolites or ezetimibe/metabolites are excreted in human milk. Because of the potential for serious adverse reactions, women taking Bempedoic acid + ezetimibe should not breastfeed their infants. Bempedoic acid + ezetimibe is contraindicated during breast-feeding.

Fertility

No data on the effect of Bempedoic acid + ezetimibe on human fertility are available. Based on animal studies, no effect on reproduction or fertility is expected with Bempedoic acid + ezetimibe.

4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

Bempedoic acid + ezetimibe has minor influence on the ability to drive and use machines. When driving vehicles or using machines, it should be taken into account that dizziness has been reported with bempedoic acid and ezetimibe.

4.8 UNDESIRABLE EFFECTS

Summary of the safety profile The most commonly reported adverse reactions are hyperuricaemia (4.7%) and constipation (4.7%). In pooled placebo-controlled clinical trials with bempedoic acid, more patients on bempedoic acid compared to placebo discontinued treatment due to muscle spasms (0.7% versus 0.3%), diarrhoea (0.5% versus < 0.1%), pain in extremity (0.4% versus 0), and nausea (0.3% versus 0.2%) although differences between bempedoic acid and placebo were not significant.

System organ class (SOC)	Adverse reactions	Frequency categories
Adverse reactions with Nustendi		
Blood and lymphatic system disorders	Anaemia Decreased haemoglobin	Common
Metabolism and nutrition disorders	Hyperuricaemia ^a	Common
	Decreased appetite	Common
Nervous system disorders	Dizziness Headache	Common
Vascular disorders	Hypertension	Common
Respiratory, thoracic and mediastinal disorders	Cough	Common
Gastrointestinal disorders	Constipation Diarrhoea Abdominal pain Nausea Dry mouth Flatulence Gastritis	Common
Hepatobiliary disorders	Liver function test increased ^b	Common
Musculoskeletal and connective tissue disorders	Back pain Muscle spasms Myalgia Pain in extremity Arthralgia	Common
Renal and urinary disorders	Blood creatinine increased	Common
General disorders and administration site conditions	Fatigue Asthenia	Common

Additional adverse reactions with bempedoic acid		
Metabolism and nutrition disorders	Gout	Common
Hepatobiliary disorders	Aspartate aminotransferase increased	Common
	Alanine aminotransferase increased	Uncommon
Renal and urinary disorders	Blood urea increased Glomerular filtration rate decreased	Uncommon
Additional adverse reactions with ezetimibe		
Blood and lymphatic system disorders	Thrombocytopenia	Not known
Immune system disorders	Hypersensitivity, including rash, urticaria, anaphylaxis and angio-oedema	Not known
Psychiatric disorders	Depression	Not known
Nervous system disorders	Paraesthesia ^c	Not known
Vascular disorders	Hot flush	Uncommon
Respiratory, thoracic and mediastinal disorders	Dyspnoea	Not known

System organ class (SOC)	Adverse reactions	Frequency categories
Gastrointestinal disorders	Dyspepsia Gastrooesophageal reflux disease	Uncommon
	Pancreatitis	Not known
Hepatobiliary disorders	Aspartate aminotransferase increased Alanine aminotransferase increased Gammaglutamyltransferase increased	Uncommon
	Hepatitis Cholelithiasis Cholecystitis	Not known
Skin and subcutaneous tissue disorders	Pruritus ^c	Uncommon
	Erythema multiform	Not known
Musculoskeletal and connective tissue disorders	Blood CPK increased	Common
	Neck pain Muscular weakness ^c	Uncommon
	Myopathy/rhabdomyolysis	Not known
General disorders and administration site conditions	Chest pain Pain Oedema peripheral ^c	Uncommon

a. Hyperuricaemia includes hyperuricaemia and increased uric acid levels

b. Liver function test includes increased liver function tests and liver function tests that are abnormal

c. Adverse reactions with ezetimibe coadministered with a statin

Description of selected adverse reactions

Increased serum uric acid

Bempedoic acid + ezetimibe increases serum uric acid possibly due to inhibition of renal tubular OAT2 by bempedoic acid. A mean increase of 0.6 mg/dL (35.7 micromole/L) in uric acid compared to baseline was observed with Bempedoic acid + ezetimibe at week 12. The elevations in serum uric acid usually occurred within the first 4 weeks of treatment and returned to baseline following discontinuation of treatment. There were no reports of gout with Bempedoic acid + ezetimibe. In the pooled placebo-controlled trials of bempedoic acid, gout was reported in 1.4% of patients treated with bempedoic acid and 0.4% of patients treated with placebo. In both treatment groups, patients who reported gout were more likely to have a medical history of gout and/or baseline levels of uric acid above the ULN.

Effects on serum creatinine and blood urea nitrogen

Bempedoic acid + ezetimibe increases serum creatinine and BUN. A mean increase of 0.02 mg/dL (1.8 micromole/L) in serum creatinine and a mean increase of 2.7 mg/dL (1.0 mmol/L) in BUN compared to baseline was observed with Bempedoic acid + ezetimibe at week 12. The elevations in serum creatinine and BUN usually occurred within the first 4 weeks of treatment, remained stable, and returned to baseline following discontinuation of therapy.

The observed elevations in serum creatinine may be associated with bempedoic acid inhibition of OAT2-dependent renal tubular secretion of creatinine, representing a drug-endogenous substrate interaction, and does not appear to indicate worsening renal function. This effect should be considered when interpreting changes in estimated creatinine clearance in patients on Bempedoic acid + ezetimibe therapy, particularly in patients with medical conditions or receiving medicinal products that require monitoring of estimated creatinine clearance.

Hepatic enzyme elevations

Hepatic transaminase (AST and/or ALT) elevations of $\geq 3 \times$ ULN were reported in 2.4% of patients treated with Bempedoic acid + ezetimibe compared with no patients on placebo. In four controlled clinical trials of bempedoic acid, the incidence of elevations ($\geq 3 \times$ ULN) in hepatic transaminase levels (AST and/or ALT) was 0.7% for patients treated with bempedoic acid and 0.3% for placebo. In controlled clinical combination trials of ezetimibe initiated concurrently with a statin, the incidence of consecutive elevations ($\geq 3 \times$ ULN) in hepatic transaminase levels was 1.3% for patients treated with ezetimibe administered with statins and 0.4% for patients treated with statins alone. The elevations in transaminases with bempedoic acid or ezetimibe were not associated with other evidence of liver dysfunction.

Decreased haemoglobin

In the pooled placebo-controlled trials of bempedoic acid, a decrease in haemoglobin from baseline of ≥ 20 g/L and $<$ lower limit of normal (LLN) was observed in 4.6% of patients in the bempedoic acid group compared with 1.9% of patients on placebo. Greater than 50 g/L and $<$ LLN decreases in haemoglobin were reported at similar rates in bempedoic acid and placebo groups (0.2% versus 0.2%, respectively). The decreases in haemoglobin usually occurred within the first 4 weeks of treatment and returned to baseline following discontinuation of treatment. Among patients who had normal haemoglobin values at baseline, 1.4% in the bempedoic acid group and 0.4% in the placebo group experienced haemoglobin values below

LLN while on treatment. Anaemia was reported in 2.5% of patients treated with bempedoic acid and 1.6% of patients treated with placebo

4.9 OVERDOSE

In the event of overdose, the patient should be treated symptomatically, and supportive measures instituted as required.

Bempedoic acid Doses up to 240 mg/day (1.3 times the approved recommended dose) have been administered in clinical trials with no evidence of dose limiting toxicity. No adverse events were observed in animal studies at exposures up to 14-fold higher than those in patients treated with bempedoic acid at 180 mg once daily.

Ezetimibe In clinical studies, administration of ezetimibe, 50 mg/day to 15 healthy subjects for up to 14 days, or 40 mg/day to 18 patients with primary hypercholesterolaemia for up to 56 days, did not result in an increase in the rate of adverse events. In animals, no toxicity was observed after single oral doses of 5000 mg/kg of ezetimibe in rats and mice and 3000 mg/kg in dogs.

5 PHARMACOLOGICAL PROPERTIES

5.1 MECHANISM OF ACTION

Bempedoic acid + ezetimibe contains bempedoic acid and ezetimibe, two LDL-C lowering compounds with complementary mechanisms of action. Bempedoic acid + ezetimibe reduces elevated LDL-C through dual inhibition of cholesterol synthesis in the liver and cholesterol absorption in the intestine.

Bempedoic acid

Bempedoic acid is an adenosine triphosphate-citrate lyase (ACL) inhibitor that lowers LDL-C by inhibition of cholesterol synthesis in the liver. ACL is an enzyme upstream of 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase in the cholesterol biosynthesis pathway. Bempedoic acid requires coenzyme A (CoA) activation by very long-chain acyl-CoA synthetase 1 (ACSVL1) to ETC-1002-CoA. ACSVL1 is expressed primarily in the liver and not in skeletal muscle. Inhibition of ACL by ETC-1002-CoA results in decreased cholesterol synthesis in the liver and lowers LDL-C in blood via upregulation of low-density lipoprotein receptors. Additionally, inhibition of ACL by ETC-1002-CoA results in concomitant suppression of hepatic fatty acid biosynthesis.

Ezetimibe

Ezetimibe reduces blood cholesterol by inhibiting the absorption of cholesterol by the small intestine. The molecular target of ezetimibe has been shown to be the sterol transporter, Niemann-Pick C1-Like 1 (NPC1L1), which is involved in the intestinal uptake of cholesterol and phytosterols. Ezetimibe localizes at the brush border of the small intestine and inhibits the absorption of cholesterol, leading to a decrease in the delivery of intestinal cholesterol to the liver.

5.2 PHARMACODYNAMIC PROPERTIES

Administration of bempedoic acid and ezetimibe alone and in combination with other lipid modifying medicinal products decreases LDL-C, non-high density lipoprotein cholesterol

(non-HDL-C), apolipoprotein B (apo B), and total cholesterol (TC) in patients with hypercholesterolaemia or mixed dyslipidaemia.

Because patients with diabetes are at elevated risk for atherosclerotic cardiovascular disease, the clinical trials of bempedoic acid included patients with diabetes mellitus. Among the subset of patients with diabetes, lower levels of HbA1c were observed as compared to placebo (on average 0.2%). In patients without diabetes, no difference in HbA1c was observed between bempedoic acid and placebo and there were no differences in the rates of hypoglycaemia.

Cardiac electrophysiology A QT trial has been conducted for bempedoic acid. At a dose of 240 mg (1.3 times the approved recommended dose), bempedoic acid does not prolong the QT interval to any clinically relevant extent. The effect of ezetimibe or the combination regimen Bempedoic acid + ezetimibe on QT interval has not been evaluated.

5.3 PHARMACOKINETIC PROPERTIES

Absorption

Bempedoic acid + ezetimibe

The bioavailability of Bempedoic acid + ezetimibe tablets was similar relative to that from the individual tablets, coadministered. C_{max} values for bempedoic acid and its active metabolite (ESP15228) were similar between formulations, but ezetimibe and ezetimibe glucuronide C_{max} values were approximately 13% and 22% lower, respectively, for Bempedoic acid + ezetimibe relative to the individual tablets, coadministered. Given a similar overall extent of ezetimibe and ezetimibe glucuronide exposure (as measured by AUC), a 22% lower C_{max} is unlikely to be clinically significant.

No clinically significant pharmacokinetic interaction was seen when ezetimibe was coadministered with bempedoic acid. Total ezetimibe (ezetimibe and its glucuronide form) and ezetimibe glucuronide AUC and C_{max} increased approximately 1.6- and 1.8-fold, respectively, when a single dose of ezetimibe was taken with steady-state bempedoic acid. This increase is likely due to inhibition of OATP1B1 by bempedoic acid, which results in decreased hepatic uptake and subsequently decreased elimination of ezetimibe-glucuronide. Increases in the AUC and C_{max} for ezetimibe were less than 20%.

Bempedoic acid

Pharmacokinetic data indicate that bempedoic acid is absorbed with a median time to maximum concentration of 3.5 hours when administered as Bempedoic acid + ezetimibe 180 mg tablets. Bempedoic acid pharmacokinetic parameters are presented as the mean [standard deviation (SD)] unless otherwise specified. Bempedoic acid can be considered a prodrug that is activated intracellularly by ACSVL1 to ETC-1002-CoA. The steady-state C_{max} and AUC following multiple dose administration in patients with hypercholesterolaemia were 24.8 (6.9) microgram/mL and 348 (120) microgram·h/mL, respectively. Bempedoic acid steady-state pharmacokinetics were generally linear over a range of 120 mg to 220 mg. There were no time-dependent changes in bempedoic acid pharmacokinetics following repeat administration at the recommended dose, and bempedoic acid steady-state was achieved after 7 days. The mean accumulation ratio of bempedoic acid was approximately 2.3-fold.

Ezetimibe

After oral administration, ezetimibe is rapidly absorbed and extensively conjugated to a pharmacologically active phenolic glucuronide (ezetimibe-glucuronide). Mean C_{max} occur within 1 to 2 hours for ezetimibe-glucuronide and 4 to 12 hours for ezetimibe. The absolute

bioavailability of ezetimibe cannot be determined as the compound is virtually insoluble in aqueous media suitable for injection. Ezetimibe undergoes extensive enterohepatic cycling, multiple peaks of ezetimibe can be observed.

Effect of food

After the administration of Bempedoic acid + ezetimibe with a high-fat, high calorie breakfast in healthy subjects, the AUC for bempedoic acid and ezetimibe were comparable to the fasted state. Compared to the fasted state, the fed state resulted in 30% and 12% reductions in C_{max} of bempedoic acid and ezetimibe, respectively. Relative to the fasted state, the fed state resulted in 12% and 42% reductions in ezetimibe glucuronide AUC and C_{max}, respectively. This effect of food is not considered to be clinically meaningful.

Distribution

Bempedoic acid

The bempedoic acid apparent volume of distribution (V/F) was 18 L. Plasma protein binding of bempedoic acid, its glucuronide and its active metabolite, ESP15228, were 99.3%, 98.8% and 99.2%, respectively. Bempedoic acid does not partition into red blood cells.

Ezetimibe

Ezetimibe and ezetimibe-glucuronide are bound 99.7% and 88% to 92% to human plasma proteins, respectively.

Biotransformation

Bempedoic acid

In vitro metabolic interaction studies suggest that bempedoic acid, as well as its active metabolite and glucuronide forms are not metabolised by and do not inhibit or induce cytochrome P450 enzymes.

The primary route of elimination for bempedoic acid is through metabolism to the acyl glucuronide. Bempedoic acid is also reversibly converted to an active metabolite (ESP15228) based on aldo-keto reductase activity observed in vitro from human liver. Mean plasma AUC metabolite/parent drug ratio for ESP15228 following repeat-dose administration was 18% and remained constant over time. Both bempedoic acid and ESP15228 are converted to inactive glucuronide conjugates in vitro by UGT2B7. Bempedoic acid, ESP15228 and their respective conjugated forms were detected in plasma with bempedoic acid accounting for the majority (46%) of the AUC_{0-48h} and its glucuronide being the next most prevalent (30%). ESP15228 and its glucuronide represented 10% and 11% of the plasma AUC_{0-48h}, respectively.

The steady-state C_{max} and AUC of the equipotent active metabolite (ESP15228) of bempedoic acid in patients with hypercholesterolaemia were 3.0 (1.4) microgram/mL and 54.1 (26.4) microgram·h/mL, respectively. ESP15228 likely made a minor contribution to the overall clinical activity of bempedoic acid based on systemic exposure and pharmacokinetic properties.

Ezetimibe

In preclinical studies, it has been shown that ezetimibe does not induce cytochrome P450 drug metabolising enzymes. No clinically significant pharmacokinetic interactions have been observed between ezetimibe and drugs known to be metabolised by cytochromes P450 1A2, 2D6, 2C8, 2C9, and 3A4, or N-acetyltransferase. Ezetimibe is metabolised primarily in the small intestine and liver via glucuronide conjugation (a phase II reaction) with subsequent biliary excretion. Minimal oxidative metabolism (a phase I reaction) has been observed in all

species evaluated. Ezetimibe and ezetimibe glucuronide are the major drug-derived compounds detected in plasma, constituting approximately 10% to 20% and 80% to 90% of the total drug in plasma, respectively. Both ezetimibe and ezetimibe glucuronide are slowly eliminated from plasma with evidence of significant enterohepatic recycling.

Elimination

Bempedoic acid The steady-state clearance (CL/F) of bempedoic acid determined from a population PK analysis in patients with hypercholesterolaemia was 12.1 mL/min after once-daily dosing; renal clearance of unchanged bempedoic acid represented less than 2% of total clearance. The mean (SD) half-life for bempedoic acid in humans was 19 (10) hours at steady-state.

Following single oral administration of 240 mg of bempedoic acid (1.3 times the approved recommended dose), 62.1% of the total dose (bempedoic acid and its metabolites) was recovered in urine, primarily as the acyl glucuronide conjugate of bempedoic acid, and 25.4% was recovered in faeces. Less than 5% of the administered dose was excreted as unchanged bempedoic acid in faeces and urine combined.

Ezetimibe

Following oral administration of ¹⁴C-ezetimibe (20 mg) to human subjects, total ezetimibe (ezetimibe and ezetimibe-glucuronide) accounted for approximately 93% of the total radioactivity in plasma. Approximately 78% and 11% of the administered radioactivity were recovered in the faeces and urine, respectively, over a 10-day collection period. After 48 hours, there were no detectable levels of radioactivity in the plasma. The half-life for ezetimibe and ezetimibe-glucuronide is approximately 22 hours.

Special populations

Renal impairment

Bempedoic acid

Pharmacokinetics of bempedoic acid was evaluated in a population PK analysis performed on pooled data from all clinical trials (n=2,261) to assess renal function on the steady-state AUC of bempedoic acid and in a single-dose pharmacokinetic study in subjects with varying degrees of renal function. Compared to patients with normal renal function, the mean bempedoic acid exposures were higher in patients with mild or moderate renal impairment by 1.4-fold (90% PI: 1.3, 1.4) and 1.9-fold (90% PI: 1.7, 2.0), respectively.

There is limited information in patients with severe renal impairment; in a single dose study, the bempedoic acid AUC was increased by 2.4-fold in patients (n=5) with severe renal impairment (eGFR < 30 mL/min/1.73 m²) compared to those with normal renal function. Clinical studies of Bempedoic acid + ezetimibe did not include patients with ESRD on dialysis.

Ezetimibe

After a single 10 mg dose of ezetimibe in patients with severe renal disease (n=8; mean CrCl ≤ 30 mL/min/1.73 m²), the mean AUC for total ezetimibe was increased approximately 1.5-fold, compared to healthy subjects (n=9). This result is not considered clinically significant. An additional patient in this study (post-renal transplant and receiving multiple medicinal products, including ciclosporin) had a 12-fold greater exposure to total ezetimibe.

Hepatic impairment Bempedoic acid + ezetimibe is not recommended in patients with moderate or severe hepatic impairment due to the unknown effects of increased exposure to ezetimibe.

Bempedoic acid

The pharmacokinetics of bempedoic acid and its metabolite (ESP15228) was studied in patients with normal hepatic function or mild or moderate hepatic impairment (Child-Pugh A or B) following a single dose (n=8/group). Compared to patients with normal hepatic function, the bempedoic acid mean C_{max} and AUC were decreased by 11% and 22%, respectively, in patients with mild hepatic impairment and by 14% and 16%, respectively, in patients with moderate hepatic impairment. This is not expected to result in lower efficacy. Bempedoic acid was not studied in patients with severe hepatic impairment (Child-Pugh C).

Ezetimibe

After a single 10 mg dose of ezetimibe, the mean AUC for total ezetimibe was increased approximately 1.7-fold in patients with mild hepatic impairment (Child-Pugh A), compared with healthy subjects. In a 14-day, multiple-dose study (10 mg daily) in patients with moderate hepatic impairment (Child-Pugh B), the mean AUC for total ezetimibe was increased approximately 4-fold on Day 1 and Day 14 compared with healthy subjects.

Other special populations

Bempedoic acid

Of the 3,621 patients treated with bempedoic acid in the placebo-controlled studies, 2,098 (58%) were > 65 years old. No overall differences in safety or efficacy were observed between these patients and younger patients.

The pharmacokinetics of bempedoic acid were not affected by age, gender, or race. Body weight was a statistically significant covariate. The lowest quartile of body weight (< 73 kg) was associated with an approximate 30% greater exposure. The increase in exposure was not clinically significant and no dose adjustments are recommended based on weight.

Ezetimibe

Geriatrics

In a multiple-dose study with ezetimibe given 10 mg once daily for 10 days, plasma concentrations for total ezetimibe were about 2-fold higher in older (≥ 65 years) healthy subjects compared to younger subjects. LDL-C reduction and safety profile are comparable between elderly and young subjects treated with ezetimibe.

Gender

Plasma concentrations for total ezetimibe are slightly higher (approximately 20%) in women than in men. LDL-C reduction and safety profile are comparable between men and women treated with ezetimibe.

6. NONCLINICAL PROPERTIES

6.1 ANIMAL TOXICOLOGY OR PHARMACOLOGY

Bempedoic acid + ezetimibe

Coadministration of bempedoic acid with doses of ezetimibe in rats at systemic total exposures > 50 times the human clinical exposure did not alter the toxicologic profile of either bempedoic acid or ezetimibe. Bempedoic acid in combination with ezetimibe did not alter the effects on embryo-fetal development profile of bempedoic acid or ezetimibe.

Bempedoic acid

The standard battery of genotoxicity studies have not identified any mutagenic or clastogenic potential of bempedoic acid. In full lifetime carcinogenicity studies in rodents, bempedoic acid increased the incidence of hepatocellular and thyroid gland follicular tumours in male rats and hepatocellular tumours in male mice. Because these are common tumours observed in rodent lifetime bioassays and the mechanism for tumourigenesis is secondary to a rodent-specific PPAR alpha activation, these tumours are not considered to translate to human risk.

Increased liver weight and hepatocellular hypertrophy were observed in rats only and were partially reversed after the 1-month recovery at ≥ 30 mg/kg/day or 4 times the exposure in humans at 180 mg. Reversible, non-adverse changes in laboratory parameters indicative of these hepatic effects, decreases in red blood cell and coagulation parameters, and increases in urea nitrogen and creatinine were observed in both species at tolerated doses. The NOAEL for adverse response in the chronic studies was 10 mg/kg/day and 60 mg/kg/day associated with exposures below and 15 times the human exposure at 180 mg in rats and monkeys, respectively.

Bempedoic acid was not teratogenic or toxic to embryos or foetuses in pregnant rabbits at doses up to 80 mg/kg/day or 12 times the systemic exposure in humans at 180 mg. Pregnant rats given bempedoic acid at 10, 30, and 60 mg/kg/day during organogenesis had decreased numbers of viable foetuses and reduced foetal body weight at ≥ 30 mg/kg/day or 4 times the systemic exposure in humans at 180 mg. An increased incidence of foetal skeletal findings (bent scapula and ribs) were observed at all doses, at exposures below the systemic exposure in humans at 180 mg. In a pre- and post-natal development study, pregnant rats administered bempedoic acid at 5, 10, 20 and 30 mg/kg/day throughout pregnancy and lactation had adverse maternal effects at ≥ 20 mg/kg/day and reductions in numbers of live pups and pup survival, pup growth and learning and memory at ≥ 10 mg/kg/day, with maternal exposures at 10 mg/kg/day, less than the exposure in humans at 180 mg.

Administration of bempedoic acid to male and female rats prior to mating and through gestation day 7 in females resulted in changes in estrous cyclicity, decreased numbers of corpora lutea and implants at ≥ 30 mg/kg/day with no effects on male or female fertility or sperm parameters at 60 mg/kg/day (4 and 9 times the systemic exposure in humans at 180 mg, respectively).

Ezetimibe

Animal studies on the chronic toxicity of ezetimibe identified no target organs for toxic effects. In dogs treated for four weeks with ezetimibe (≥ 0.03 mg/kg/day) the cholesterol concentration in the cystic bile was increased by a factor of 2.5 to 3.5. However, in a one-year study in dogs given doses of up to 300 mg/kg/day no increased incidence of cholelithiasis or other hepatobiliary effects were observed. The significance of these data for humans is not known. A lithogenic risk associated with the therapeutic use of ezetimibe cannot be ruled out.

In coadministration studies with ezetimibe and statins the toxic effects observed were essentially those typically associated with statins. Some of the toxic effects were more pronounced than observed during treatment with statins alone. This is attributed to pharmacokinetic and pharmacodynamic interactions in coadministration therapy. Myopathies

occurred in rats only after exposure to doses that were several times higher than the human therapeutic dose (approximately 20 times the AUC level for statins and 500 to 2,000 times the AUC level for the active metabolites).

In a series of in vivo and in vitro assays ezetimibe, given alone or coadministered with statins, exhibited no genotoxic potential. Long-term carcinogenicity tests on ezetimibe were negative.

Ezetimibe had no effect on the fertility of male or female rats, nor was it found to be teratogenic in rats or rabbits, nor did it affect prenatal or postnatal development. Ezetimibe crossed the placental barrier in pregnant rats and rabbits given multiple doses of 1,000 mg/kg/day. The coadministration of ezetimibe and statins was not teratogenic in rats. In pregnant rabbits a small number of skeletal deformities (fused thoracic and caudal vertebrae, reduced number of caudal vertebrae) were observed. The coadministration of ezetimibe with lovastatin resulted in embryolethal effects

7. DESCRIPTION

Bemneo EZ is supplied for oral administration, as a film coated tablet of Bempedoic Acid and Ezetimibe. Each film coated tablet of Bemneo EZ contains Bempedoic Acid 180mg and Ezetimibe 10mg.

8. PHARMACEUTICAL PARTICULARS

8.1 INCOMPATIBILITIES

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

8.2 SHELF-LIFE

Refer pack

8.3 Packaging Information

Refer pack

8.4 STORAGE CONDITION AND HANDING INSTRUCTIONS

Refer pack

9. PATIENT COUNSELLING INFORMATION

Advise patients to read the patient labelling Risk of Hyperuricemia

Advise patients of the risk of elevated serum uric acid levels, including development of gout. Inform patients that serum uric acid levels may be monitored during treatment with Bempedoic acid and Ezetimibe. Patients with signs or symptoms of hyperuricemia should contact their healthcare provider if symptoms occur. [see warning and precautions]

Risk of Tendon Rupture

Inform patients of the risk of tendon rupture. Advise patients to rest at the first sign of tendinitis or tendon rupture and to immediately contact their healthcare provider if tendinitis or tendon rupture symptoms occur. [see warning and precautions]

Risk of Myopathy with Concomitant Use of Simvastatin or Pravastatin

Advise patients to notify their healthcare provider(s) if they are taking, or plan to take simvastatin or pravastatin. The risk of myopathy occurring with the use of simvastatin or pravastatin may be increased when taken with Bempedoic acid and Ezetimibe Tablet.

Version: 2.0, Dated: 22nd Feb 2024

Replaces Version: 1.0, Dated: 02nd Feb 2023

Pregnancy

Advise pregnant women of the potential risk to a fetus based on Bempedoic acid and Ezetimibe mechanism of action. Advise females to inform their healthcare provider of a known or suspected pregnancy.

10. DETAILS OF MANUFACTURER

Refer Pack for manufacturer details.

Marketed By:

Abbott Healthcare Pvt. Ltd.

Angel Space, Bldg. D-4, Gala No. 1 to 6 &

11 to 16 Ground Floor, 101 to 106 &

111 to 116 First Floor, 201 to 206 & 211 to 216

2nd Floor, Pimpalas, Dist. Thane, Bhiwandi - 421 302, India.

11. DETAILS OF PERMISSION OR LICENCE NUMBER

Refer Pack for Permission/License details.

12. DATE OF REVISION

Version 2.0, Dated: 22nd Feb 2024

For Product Complaints/Adverse events or Queries please write to
webmasterindia@abbott.com