

For the use of a Registered Medical Practitioner (neurologist only)

1. GENERIC NAME AND BRAND NAME

Oxcarbazepine Tablets I.P. 150mg

Oxmazetol® 150

Oxcarbazepine Tablets I.P. 300mg

Oxmazetol® 300

Oxcarbazepine Tablets I.P. 450mg

Oxmazetol® 450

Oxcarbazepine Tablets I.P. 600mg

Oxmazetol® 600

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Oxcarbazepine Tablets I.P. 150mg

Each film coated tablet contains:

Oxcarbazepine I.P. 150mg

Excipients q.s.

Colour: Lake of Tartrazine, Brilliant Blue FCF & Titanium Dioxide I.P.

Oxcarbazepine Tablets I.P. 300mg

Each film coated tablet contains:

Oxcarbazepine I.P. 300mg

Excipients q.s.

Colour: Lake of Ponceau 4R & Titanium Dioxide I.P.

Oxcarbazepine Tablets I.P. 450mg

Each film coated tablet contains:

Oxcarbazepine I.P. 450 mg

Excipients q.s.

Colour: Lake of Sunset Yellow FCF & Titanium Dioxide I.P.

Oxcarbazepine Tablets I.P. 600mg

Each film coated tablet contains:

Oxcarbazepine I.P. 600mg

Excipients q.s.

Colour: Lake of Brilliant Blue FCF & Titanium Dioxide I.P.

3. DOSAGE FORM AND STRENGTH

Refer section 1 & 2

4. CLINICAL PARTICULARS

4.1 THERAPEUTIC INDICATION

- Monotherapy in the treatment of partial seizures in children ages 4- 16.
- Treatment of generalized tonic clonic seizures in adults and children.

Version 5.0; Dated 4-May-26

Replaces 4.0; Dated 11th Sep 2023

- For additional indication of adjunctive therapy in the treatment of partial seizures in children in the age group of 4-14 years with epilepsy

4.2 POSOLOGY AND METHOD OF ADMINISTRATION

Posology:

In mono- and adjunctive therapy, treatment with Oxcarbazepine is initiated with a clinically effective dose given in two divided doses. The dose may be increased depending on the clinical response of the patient. When other antiepileptic medicinal products are replaced by Oxcarbazepine, the dose of the concomitant antiepileptic medicinal product(s) should be reduced gradually on initiation of Oxcarbazepine therapy. In adjunctive therapy, as the total antiepileptic medicinal product load of the patient is increased, the dose of concomitant antiepileptic medicinal product(s) may need to be reduced and/or the Oxcarbazepine dose increased more slowly.

Therapeutic drug monitoring

The therapeutic effect of oxcarbazepine is primarily exerted through the active metabolite 10-monohydroxy derivative (MHD) of oxcarbazepine.

Plasma level monitoring of oxcarbazepine or MHD is not routinely warranted. However, may be useful in situations where an alteration in MHD clearance is to be expected. In such situations, the dose of oxcarbazepine may be adjusted (based on plasma levels measured 2-4 hours post dose) to maintain peak MHD plasma levels < 35 mg/L.

Adults

Monotherapy

Recommended initial dose

Oxcarbazepine should be initiated with a dose of 600 mg/day (8-10 mg/kg/day) given in 2 divided doses.

Maintenance dose

If clinically indicated, the dose may be increased by a maximum of 600 mg/day at approximately weekly intervals from the starting dose to achieve the desired clinical response. Therapeutic effects are seen at doses between 600 mg/day and 2400 mg/day.

Controlled monotherapy trials in patients not currently being treated with antiepileptic medicinal products showed 1200 mg/day to be an effective dose; however, a dose of 2400 mg/day has been shown to be effective in more refractory patients converted from other antiepileptic medicinal products to oxcarbazepine monotherapy.

Maximum recommended dose

In a controlled hospital setting, dose increases up to 2400 mg/day have been achieved over 48 hours.

Adjunctive therapy

Recommended initial dose

Oxcarbazepine should be initiated with a dose of 600 mg/day (8-10 mg/kg/day) given in 2 divided doses.

Maintenance dose

If clinically indicated, the dose may be increased by a maximum of 600 mg/day at approximately weekly intervals from the starting dose to achieve the desired clinical response. Therapeutic response are seen at doses between 600 mg/day and 2400 mg/day.

Maximum recommended dose

Daily doses from 600 to 2400 mg/day have been shown to be effective in a controlled adjunctive therapy trial, although most patients were not able to tolerate the 2400 mg/day dose without reduction of concomitant antiepileptic medicinal products, mainly because of CNS-related adverse events.

Daily doses above 2400 mg/day have not been studied systematically in clinical trials.

Elderly (65 years old and above)

No special dose recommendations are necessary in elderly patients because therapeutic doses are individually adjusted. Dosage adjustments are recommended in elderly patients with renal impairment (creatinine clearance <30 ml/min) (see 'Patients with renal impairment'). Close monitoring of sodium levels is required inpatients at risk of hyponatraemia.

Patients with hepatic impairment

No dosage adjustment is required for patients with mild to moderate hepatic impairment. Oxcarbazepine has not been studied in patients with severe hepatic impairment, therefore, caution should be exercised when dosing severely impaired patients.

Patients with renal impairment

In patients with impaired renal function (creatinine clearance less than 30 ml/min) oxcarbazepine therapy should be initiated at half the usual starting dose (300 mg/day) and increased, in at least weekly intervals, to achieve the desired clinical response.

Dose escalation in renally impaired patients may require more careful observation.

Paediatric population

Recommended initial dose

In mono- and adjunctive therapy, Oxcarbazepine should be initiated with a dose of 8-10 mg/kg/day given in 2 divided doses.

Maintenance dose

In adjunctive therapy trials, a maintenance dose of 30-46 mg/kg/day, achieved over two weeks, is shown to be effective and well tolerated in children. Therapeutic effects were seen at a median maintenance dose of approximately 30 mg/kg/day.

Maximum recommended dose

If clinically indicated, the dose may be increased by a maximum of 10 mg/kg/day at approximately weekly intervals from the starting dose, to a maximum dose of 46 mg/kg/day, to achieve the desired clinical response.

Oxcarbazepine is recommended for use in children of 6 years of age and above. Safety and efficacy have been evaluated in controlled clinical trials involving approximately 230 children aged less than 6 years (down to 1 month). Oxcarbazepine is not recommended in children aged less than 6 years since safety and efficacy have not been adequately demonstrated.

All the above dosing recommendations (adults, elderly and children) are based on the doses studied in clinical trials for all age groups. However, lower initiation doses may be considered where appropriate.

Method of administration

Oral use.

Oxcarbazepine can be taken with or without food.

The tablets are scored and can be broken in two halves in order to make it easier for the patient to swallow the tablet. For children, who cannot swallow tablets or where the required dose cannot be administered using tablets, other oxcarbazepine containing pharmaceutical forms are available.

4.3 CONTRAINDICATIONS

Hypersensitivity to the active substance, eslicarbazepine or to any of the other Antiepileptics drugs.

4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Hypersensitivity

Class I (immediate) hypersensitivity reactions including rash, pruritus, urticaria, angioedema and reports of anaphylaxis have been received in the post-marketing period. Cases of anaphylaxis and angioedema involving the larynx, glottis, lips and eyelids have been reported in patients after taking the first or subsequent doses of oxcarbazepine. If a patient develops these reactions after treatment with oxcarbazepine, the drug should be discontinued and an alternative treatment started.

Patients who have exhibited hypersensitivity reactions to carbamazepine should be informed that approximately 25-30 % of these patients may experience hypersensitivity reactions (e.g. severe skin reactions) with oxcarbazepine.

Hypersensitivity reactions, including multi-organ hypersensitivity reactions, may also occur in patients without a history of hypersensitivity to carbamazepine. Such reactions can affect the skin, liver, blood and lymphatic system or other organs, either individually or together in the context of a systemic reaction. In general, if signs and symptoms suggestive of hypersensitivity reactions occur Oxcarbazepine should be withdrawn immediately.

Dermatological effects

Serious dermatological reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis (Lyell's syndrome) and erythema multiforme, have been reported very rarely in association with the use of oxcarbazepine. Patients with serious dermatological reactions may require hospitalisation, as these conditions may be life-threatening and very rarely be fatal. Oxcarbazepine associated cases occurred in both children and adults.

The median time to onset was 19 days. Several isolated cases of recurrence of the serious skin reaction when rechallenged with oxcarbazepine were reported. Patients who develop a skin reaction with oxcarbazepine should be promptly evaluated and oxcarbazepine withdrawn immediately unless the rash is clearly not drug related. In case of treatment withdrawal, consideration should be given to replacing oxcarbazepine with other antiepileptic drug therapy to avoid withdrawal seizures. Oxcarbazepine should not be restarted in patients who discontinued treatment due to a hypersensitivity reaction.

Version 5.0; Dated 4-May-26

Replaces 4.0; Dated 11th Sep 2023

HLA-B*1502 allele – in Han Chinese, Thai and other Asian populations

HLA-B*1502 in individuals of Han Chinese and Thai origin has been shown to be strongly associated with the risk of developing the severe cutaneous reactions known as Stevens-Johnson syndrome (SJS)/ toxic epidermal necrolysis (TEN), when treated with carbamazepine. The chemical structure of oxcarbazepine is similar to that of carbamazepine, and it is possible that patients who are positive for HLA-B*1502 may also be at risk for SJS/TEN after treatment with oxcarbazepine. There are some data that suggest that such an association exists for oxcarbazepine. The prevalence of HLA-B*1502 carrier is about 10% in Han Chinese and Thai populations. Whenever possible, these individuals should be screened for this allele before starting treatment with carbamazepine or a chemically-related active substance. If patients of these origins are tested positive for HLA-B*1502 allele, the use of oxcarbazepine may be considered if the benefits are thought to exceed risks.

Because of the prevalence of this allele in other Asian populations (e.g. above 15% in the Philippines and Malaysia), testing genetically at risk populations for the presence of HLA-B*1502 may be considered.

The prevalence of the HLA-B*1502 allele is negligible in e.g. European descent, African, Hispanic populations sampled, and in Japanese and Koreans (< 1%).

Allele frequencies refer to the percentage of chromosomes in the population that carry a given allele. Since a person carries two copies of each chromosome, but even one copy of the HLA-B* 1502 allele may be enough to increase the risk of SJS, the percentage of patients who may be at risk is nearly twice the allele frequency.

HLA-A*3101 allele – European descent and Japanese populations

There are some data that suggest HLA-A*3101 is associated with an increased risk of carbamazepine induced cutaneous adverse drug reactions including SJS, TEN, Drug rash with eosinophilia (DRESS), or less severe acute generalised exanthematous pustulosis (AGEP) and maculopapular rash in people of European descent and the Japanese.

The frequency of the HLA-A*3101 allele varies widely between ethnic populations. HLA-A*3101 allele has a prevalence of 2 to 5% in European populations and about 10% in Japanese population.

The presence of HLA-A*3101 allele may increase the risk for carbamazepine induced cutaneous reactions (mostly less severe) from 5.0% in general population to 26.0% among subjects of European ancestry, whereas its absence may reduce the risk from 5.0% to 3.8%.

HLA-A*3101 allele- Other descents

The frequency of this allele is estimated to be less than 5% in the majority of Australian, Asian, African and North American populations with some exceptions within 5 to 12%. Frequency above 15% has been estimated in some ethnic groups in South America (Argentina and Brazil), North America (US Navajo and Sioux, and Mexico Sonora Seri) and Southern India (Tamil Nadu) and between 10% to 15% in other native ethnicities in these same regions.

Allele frequencies refer to the percentage of chromosomes in the population that carry a given allele. Since a person carries two copies of each chromosome, but even one copy of the HLA-A*3101 allele may be enough to increase the risk of SJS, the percentage of patients who may be at risk is nearly twice the allele frequency.

There are insufficient data supporting a recommendation for HLA-A*3101 screening before starting carbamazepine or chemically-related compounds treatment.

If patients of European descent or Japanese origin are known to be positive for HLA-A*3101 allele, the use of carbamazepine or chemically-related compounds may be considered if the benefits are thought to exceed risks.

Limitation of genetic screening

Genetic screening results must never substitute appropriate clinical vigilance and patient management. Many Asian patients positive for HLA-B* 1502 and treated with oxcarbazepine will not develop SJS/TEN, and patients negative for HLA- B* 1502 of any ethnicity can still develop SJS/TEN. The same is true for HLA-A*3101 with respect to risk of SJS, TEN, DRESS, AGEP or maculopapular rash. The development of these severe cutaneous adverse reactions and its related morbidity due to other possible factors such as AED dose, compliance, concomitant medications, co- morbidities, and the level of dermatologic monitoring have not been studied.

Information for healthcare professionals

If testing for the presence of the HLA-B*1502 allele is performed, high-resolution "HLA-B*1502 genotyping" is recommended. The test is positive if either one or two HLA-B* 1502 alleles are detected, and negative if no HLA- B* 1502 alleles are detected. Similarly, if testing for the presence of the HLA-A*3101 allele is performed, high resolution "HLA-A*3101 genotyping" is recommended. The test is positive if either one or two HLA-A*3101 alleles are detected, and negative if no HLA-A*3101 alleles are detected.

Risk of seizure aggravation

Risk of seizure aggravation has been reported with oxcarbazepine. The risk of seizure aggravation is seen especially in children but may also occur in adults. In case of seizure aggravation, oxcarbazepine should be discontinued.

Hyponatraemia

Serum sodium levels below 125 mmol/l, usually asymptomatic and not requiring adjustment of therapy, have been observed in up to 2.7 % of oxcarbazepine treated patients. Experience from clinical trials shows that serum sodium levels returned towards normal when the oxcarbazepine dosage was reduced, discontinued or the patient was treated conservatively (e.g. restricted fluid intake). In patients with pre-existing renal conditions associated with low sodium levels (e.g. inappropriate ADH secretion like syndrome) or in patients treated concomitantly with sodium-lowering medicinal products (e.g. diuretics, desmopressin) as well as NSAIDs (e.g. indometacin), serum sodium levels should be measured prior to initiating therapy.

Thereafter, serum sodium levels should be measured after approximately two weeks and then at monthly intervals for the first three months during therapy, or according to clinical need. These risk factors may apply especially to elderly patients. For patients on oxcarbazepine therapy when starting on sodium-lowering medicinal products, the same approach for sodium checks should be followed. In general, if clinical symptoms suggestive of hyponatraemia occur in oxcarbazepine therapy, serum sodium levels measurement may be considered. Other patients may have serum sodium assessed as part of their routine laboratory studies.

All patients with cardiac insufficiency and secondary heart failure should have regular weight measurements to determine occurrence of fluid retention. In case of fluid retention or worsening of the cardiac condition, serum sodium levels should be checked. If hyponatraemia

is observed, water restriction is an important counter-measurement. As oxcarbazepine may, very rarely, lead to impairment of cardiac conduction, patients with pre-existing conduction disturbances (e.g. atrioventricular-block, arrhythmia) should be followed carefully.

Hypothyroidism

Hypothyroidism is an adverse drug reaction of oxcarbazepine. Considering the importance of thyroid hormones in children's development after birth, thyroid function monitoring is recommended in the pediatric age group while on oxcarbazepine therapy.

Hepatic function

Very rare cases of hepatitis have been reported, which in most of the cases resolved favourably. When a hepatic event is suspected, liver function should be evaluated and discontinuation of Oxcarbazepine should be considered. Caution should be exercised when treating patients with severe hepatic impairment.

Renal function

In patients with impaired renal function (creatinine clearance less than 30 mL/min), caution should be exercised during oxcarbazepine treatment especially with regard to the starting dose and up titration of the dose. Plasma level monitoring of MHD may be considered.

Haematological effects

Very rare reports of agranulocytosis, aplastic anemia and pancytopenia have been seen in patients treated with oxcarbazepine during post-marketing experience. Discontinuation of the medicinal product should be considered if any evidence of significant bone marrow depression develops.

Suicidal behaviour

Suicidal ideation and behaviour have been reported in patients treated with antiepileptic medicinal products in several indications. A meta-analysis of randomised placebo controlled trials of antiepileptic drugs has also shown a small increased risk of suicidal ideation and behaviour. The mechanism of this risk is not known and the available data do not exclude the possibility of an increased risk for oxcarbazepine.

Therefore, patients should be monitored for signs of suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

Hormonal contraceptives

Female patients of childbearing age should be warned that the concurrent use of oxcarbazepine with hormonal contraceptives may render this type of contraceptive ineffective. Additional non-hormonal forms of contraception are recommended when using oxcarbazepine.

Alcohol

Caution should be exercised if alcohol is taken in combination with oxcarbazepine therapy, due to a possible additive sedative effect.

Withdrawal

As with all antiepileptic medicinal products, oxcarbazepine should be withdrawn gradually to minimise the potential of increased seizure frequency.

Monitoring of plasma levels

Although correlations between dosage and plasma levels of oxcarbazepine, and between plasma levels and clinical efficacy or tolerability are rather tenuous, monitoring of the plasma levels may be useful in the following situations in order to rule out noncompliance or in situations where an alteration in MHD clearance is to be expected, including:

- changes in renal function (see renal impairment).
- pregnancy.
- concomitant use of liver enzyme-inducing drugs.

Excipients

This medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

4.5 DRUGS INTERACTIONS

Enzyme induction

Oxcarbazepine and its pharmacologically active metabolite (the monohydroxy derivative, MHD) are weak inducers in vitro and in vivo of the cytochrome P450 enzymes CYP3A4 and CYP3A5 responsible for the metabolism of a very large number of medicinal products, for example, immunosuppressants (e.g. ciclosporin, tacrolimus), oral contraceptives (see below), and some other antiepileptic medicinal products (e.g. carbamazepine) resulting in a lower plasma concentration of these medicinal products (see table below summarising results with other antiepileptic medicinal products).

In vitro, oxcarbazepine and MHD are weak inducers of UDP-glucuronyl transferases (effects on specific enzymes in this family are not known). Therefore, in vivo oxcarbazepine and MHD may have a small inducing effect on the metabolism of medicinal products which are mainly eliminated by conjugation through the UDP-glucuronyl transferases. When initiating treatment with oxcarbazepine or changing the dose, it may take 2 to 3 weeks to reach the new level of induction.

In case of discontinuation of oxcarbazepine therapy, a dose reduction of the concomitant medications may be necessary and should be decided upon by clinical and/or plasma level monitoring. The induction is likely to gradually decrease over 2 to 3 weeks after discontinuation.

Hormonal contraceptives: Oxcarbazepine was shown to have an influence on the two components, ethinylestradiol (EE) and levonorgestrel (LNG), of an oral contraceptive. The mean AUC values of EE and LNG were decreased by 48-52 % and 32-52% respectively.

Therefore, concurrent use of oxcarbazepine with hormonal contraceptives may render these contraceptives ineffective. Another reliable contraceptive method should be used.

Enzyme inhibition

Oxcarbazepine and MHD inhibit CYP2C19. Therefore, interactions could arise when co-administering high doses of oxcarbazepine with medicinal products that are mainly metabolised by CYP2C19 (e.g. phenytoin). Phenytoin plasma levels increased by up to 40 % when oxcarbazepine was given at doses above 1,200 mg/day (see table below summarising

results with other anticonvulsants). In this case, a reduction of co-administered phenytoin may be required.

Antiepileptic and enzyme inducing medicinal products

Potential interactions between oxcarbazepine and other antiepileptic medicinal products were assessed in clinical studies. The effect of these interactions on mean AUCs and C_{min} are summarised in the following table.

Summary of antiepileptic medicinal product interactions with oxcarbazepine.

Antiepileptic medicinal product	Influence of oxcarbazepine on antiepileptic medicinal product	Influence of antiepileptic medicinal product on MHD
Co-administered	Concentration	Concentration
Carbamazepine	0 - 22 % decrease (30 % increase of carbamazepine-epoxide)	40 % decrease
Clobazam	Not studied	No influence
Felbamate	Not studied	No influence
Lamotrigine	No influence	No influence
Phenobarbital	14 - 15 % increase	30 - 31 % decrease
Phenytoin	0 - 40 % increase	29 - 35 % decrease
Valproic acid	No influence	0 - 18 % decrease

Strong inducers of cytochrome P450 enzymes and/or UGT (i.e. rifampicin, carbamazepine, phenytoin and phenobarbital) have been shown to decrease the plasma/serum levels of MHD (29-49%) in adults; in children 4 to 12 years of age, MHD clearance increased by approximately 35% when given one of the three enzyme-inducing antiepileptic medicinal products compared to monotherapy. Concomitant therapy of oxcarbazepine and lamotrigine has been associated with an increased risk of adverse events (nausea, somnolence, dizziness and headache).

When one or several antiepileptic medicinal products are concurrently administered with oxcarbazepine, a careful dose adjustment and/or plasma level monitoring may be considered on a case by case basis, notably in paediatric patients treated concomitantly with lamotrigine.

No autoinduction has been observed with oxcarbazepine.

Other medicinal product interactions

Cimetidine, erythromycin, viloxazine, warfarin and dextropropoxyphene had no effect on the pharmacokinetics of MHD.

The interaction between oxcarbazepine and MAOIs is theoretically possible based on a structural relationship of oxcarbazepine to tricyclic antidepressants.

Patients on tricyclic antidepressant therapy were included in clinical trials and no clinically relevant interactions have been observed.

The combination of lithium and oxcarbazepine might cause enhanced neurotoxicity.

4.6 USE IN SPECIAL POPULATIONS (SUCH AS PREGNANT WOMEN, LACTATING WOMEN, PAEDIATRIC PATIENTS, GERIATRIC PATIENTS ETC.)

Women of child-bearing potential and contraceptive measures

Oxcarbazepine may result in a failure of the therapeutic effect of oral contraceptive medicinal products containing ethinylestradiol (EE) and levonorgestrel (LNG). Women of child bearing potential should be advised to use highly effective contraception (preferably non-hormonal; e.g. intrauterine implants) while on treatment with oxcarbazepine.

Pregnancy

Risk related to epilepsy and antiepileptic medicinal products in general:

In the treated population, an increase in malformations has been noted with polytherapy, particularly in polytherapy including valproate.

Moreover, effective anti-epileptic therapy must not be interrupted, since the aggravation of the illness is detrimental to both the mother and the foetus.

Risk related to oxcarbazepine

There is moderate amount of data on pregnant women (300-1000 pregnancy outcomes). However, the data on oxcarbazepine associated with congenital malformation is limited. There is no increase in the total rate of malformations with oxcarbazepine as compared with the rate observed with general population (2-3%). Nevertheless, with this amount of data, a moderate teratogenic risk cannot be completely excluded.

Taking these data into consideration:

- If women receiving oxcarbazepine become pregnant or plan to become pregnant, the use of this product should be carefully re-evaluated. Minimum effective doses should be given, and monotherapy whenever possible should be preferred at least during the first three months of pregnancy.
- During pregnancy, an effective antiepileptic oxcarbazepine treatment must not be interrupted, since the aggravation of the illness is detrimental to both the mother and the foetus.

Monitoring and prevention

Some antiepileptic medicinal products may contribute to folic acid deficiency, a possible contributory cause of foetal abnormality. Folic acid supplementation is recommended before and during pregnancy. As the efficacy of this supplementation is not proved, a specific antenatal diagnosis should be offered even for women with a supplementary treatment of folic acid.

Data from a limited number of women indicate that plasma levels of the active metabolite of oxcarbazepine, the 10-monohydroxy derivative (MHD), may gradually decrease throughout pregnancy. It is recommended that clinical response should be monitored carefully in women receiving oxcarbazepine treatment during pregnancy to ensure that adequate seizure control is maintained. Determination of changes in MHD plasma concentrations should be considered. If dosages have been increased during pregnancy, postpartum MHD plasma levels may also be considered for monitoring.

In the new born child

Bleeding disorders in the new born have been reported with hepatic inductor antiepileptic medicinal products. As a precaution, vitamin K₁ should be administered as a preventive measure in the last few weeks of pregnancy and to the new born.

Version 5.0; Dated 4-May-26

Replaces 4.0; Dated 11th Sep 2023

Breastfeeding

Oxcarbazepine and its active metabolite (MHD) are excreted in human breast milk. A milk-to-plasma concentration ratio of 0.5 was found for both. The effects on the infant exposed to oxcarbazepine by this route are unknown. Therefore, oxcarbazepine should not be used during breast-feeding.

Fertility

There are no human data on fertility. In rats, oxcarbazepine had no effects on fertility. Effects on reproductive parameters in female rats were observed for MHD at doses comparable to those in humans.

4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

Adverse reactions such as dizziness, somnolence, ataxia, diplopia, blurred vision, visual disturbances, hyponatraemia and depressed level of consciousness were reported with oxcarbazepine (for complete list of ADRs), especially at the start of treatment or in connection with dose adjustments (more frequently during the up-titration phase). Patients should therefore exercise due caution when driving a vehicle or operating machinery.

4.8 UNDESIRABLE EFFECTS

Summary of the safety profile

The most commonly reported adverse reactions are somnolence, headache, dizziness, diplopia, nausea, vomiting and fatigue occurring in more than 10% of patients.

The safety profile is based on adverse events (AEs) from clinical trials assessed as related to oxcarbazepine. In addition, clinically meaningful reports on adverse experiences from named patient programs and post-marketing experience were taken into account.

Adverse drug reactions are listed by MedRA system organ class.

Tabulated list of adverse reactions

Frequency estimate*:- very common: $\geq 1/10$; common: $\geq 1/100 - < 1/10$; uncommon: $\geq 1/1,000 - < 1/100$; rare: $\geq 1/10,000 - < 1/1,000$; very rare: $< 1/10,000$; not known: cannot be estimated from the available data.

Within each system organ class, the adverse drug reactions are ranked by frequency, with the most frequent reactions first. Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Blood and lymphatic system disorders Uncommon: Very rare: Not known:	Leucopenia Thrombocytopenia Bone marrow depression, aplastic anaemia, agranulocytosis, pancytopenia, neutropenia.
Immune system disorders Very rare: Not known:	Hypersensitivity [#] Anaphylactic reactions.
Endocrine disorders Common: Not known:	Weight increased Hypothyroidism

Metabolism and nutrition disorders Common: Very rare: Not known:	Hyponatraemia [†] Hyponatraemia associated with signs and symptoms such as seizures, encephalopathy, depressed level of consciousness, confusion (see also Nervous system disorders for further undesirable effects), vision disorders (e.g. blurred vision), hypothyroidism, vomiting, nausea [†] . Inappropriate ADH secretion like syndrome with signs and symptoms of lethargy, nausea, dizziness, decrease in serum (blood) osmolality, vomiting, headache, confusional state or other neurological signs and symptoms.
Psychiatric disorders Common:	Agitation (e.g. nervousness), affect lability, confusional state, depression, apathy.
Nervous system disorders Very common: Common: Not known	Somnolence, headache, dizziness. Ataxia, tremor, nystagmus, disturbance in attention, amnesia. Speech disorders (including dysarthria); more frequent during up titration of oxcarbazepine dose
Eye disorders Very common: Common:	Diplopia. Vision blurred, visual disturbance.
Ear and labyrinth disorders Common:	Vertigo.
Cardiac disorders Very rare:	Atrioventricular block, arrhythmia.
Vascular disorders Not known:	Hypertension
Gastrointestinal disorders Very common: Common: Very rare:	Vomiting, nausea Diarrhoea, abdominal pain, constipation. Pancreatitis and/or lipase and/or amylase increase.
Hepatobiliary disorders Very rare:	Hepatitis.
Skin and subcutaneous tissue disorders Common: Uncommon: Very rare: Not known	Rash, alopecia, acne. Urticaria. Stevens-Johnson syndrome, toxic epidermal necrolysis (Lyell's syndrome), angioedema, erythema multiforme.

	Drug Rash with Eosinophilia and Systemic Symptoms (DRESS)**, Acute Generalised Exanthematous Pustulosis (AGEP)**
Musculoskeletal and connective tissue disorders Very rare: Not known	Systemic lupus erythematosus. There have been reports of decreased bone mineral density, osteopenia, osteoporosis and fractures in patients on long-term therapy with oxcarbazepine. The mechanism by which oxcarbazepine affects bone metabolism has not been identified.
General disorders and administration site conditions Very common: Common:	Fatigue. Asthenia.
Investigations Uncommon: Not known:	Hepatic enzymes increased, blood alkaline phosphatase increased. Decrease in T4 (with unclear clinical significance).
Injury, poisoning and procedural complications Not known	Fall

* According to CIOMS III frequency classification

#Hypersensitivity (including multi-organ hypersensitivity) characterised by features such as rash, fever. Other organs or systems may be affected such as blood and lymphatic system (e.g. eosinophilia, thrombocytopenia, leucopenia, lymphadenopathy, splenomegaly), liver (e.g. hepatitis, abnormal liver function tests), muscles and joints (e.g. joint swelling, myalgia, arthralgia), nervous system (e.g. hepatic encephalopathy), kidneys (e.g. renal failure, nephritis interstitial, proteinuria), lungs (e.g. pulmonary oedema, asthma, bronchospasms, interstitial lung disease, dyspnoea), angioedema.

† Serum sodium levels below 125 mmol/l have been observed in up to 2.7 % of oxcarbazepine treated patients with frequency common. In most cases, the hyponatraemia is asymptomatic and does not require adjustment of therapy. Very rarely, the hyponatraemia is associated with signs and symptoms such as seizures, encephalopathy, depressed level of consciousness, confusion, (see also Nervous system disorders for further undesirable effects), vision disorders (e.g. blurred vision), hypothyroidism, vomiting, and nausea. Low serum sodium levels generally occurred during the first 3 months of treatment with oxcarbazepine, although there were patients who first developed a serum sodium levels <125 mmol/l more than 1 year after initiation of therapy.

****Adverse drug reactions from spontaneous reports and literature cases (frequency not known):** The following adverse drug reactions have been derived from post-marketing experience with oxcarbazepine via spontaneous case reports and literature cases. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency which is therefore categorised as not known.

Paediatric population

In general, the safety profile in children was similar to that observed in the adult population.

4.9 OVERDOSE

Isolated cases of overdose have been reported. The maximum dose taken was approximately 48,000 mg.

Signs and symptoms

- Electrolyte and fluid balance conditions: hyponatraemia
- Eye disorders: diplopia, miosis, blurred vision
- Gastrointestinal disorders: nausea, vomiting, hyperkinesia
- General disorders and administration site conditions: fatigue
- Investigations: respiratory rate depression, QTc prolongation
- Nervous system disorders: drowsiness and somnolence, dizziness, ataxia and nystagmus, tremor, disturbances in coordination (coordination abnormal), convulsion, headache, coma, loss of consciousness, dyskinesia
- Psychiatric disorders: aggression, agitation, confusional state
- Vascular disorders: hypotension
- Respiratory, thoracic and mediastinal disorders: dyspnoea

Management

There is no specific antidote. Symptomatic and supportive treatment should be administered as appropriate. Removal of the medicinal product by gastric lavage and/or inactivation by administering activated charcoal should be considered.

5. PHARMACOLOGICAL PROPERTIES

5.1 MECHANISM OF ACTION

The pharmacological activity of oxcarbazepine is primarily exerted through the metabolite (MHD). The mechanism of action of oxcarbazepine and MHD is thought to be mainly based on the blockade of voltage-sensitive sodium channels, thus resulting in stabilisation of hyperexcited neural membranes, inhibition of repetitive neuronal firing, and diminishment of propagation of synaptic impulses.

In addition, increased potassium conductance and modulation of high-voltage activated calcium channels may also contribute to anticonvulsant effects. No significant interactions with brain neurotransmitter or modulator receptor sites were found.

5.2 PHARMACODYNAMIC PROPERTIES

Pharmacotherapeutic group: Antiepileptics, Carboxamide derivatives, ATC code: N03AF02

Oxcarbazepine and its active metabolite (MHD), are potent and efficacious anticonvulsants in animals. They protected rodents against generalised tonic-clonic and, to a lesser degree, clonic seizures, and abolished or reduced the frequency of chronically recurring partial seizures in Rhesus monkeys with aluminium implants. No tolerance (i.e. attenuation of anticonvulsive activity) against tonic-clonic seizures was observed when mice and rats were treated daily for 5 days or 4 weeks, respectively, with oxcarbazepine or MHD.

Pharmacodynamic/efficacy studies

A prospective, open-label, multicentre, non-comparative, 24-week observational post marketing study has been conducted in India. Out of a study population of 816 patients, 256 paediatric patients (1 month to 19 years) with generalised tonic-clonic seizures (either secondary or primary) were treated with oxcarbazepine monotherapy. The initial oxcarbazepine dose for all patients > 6 years was 8 to 10 mg/kg/day given in 2 divided doses.

For the 27 subjects aged 1 month to 6 years, the dose range for the initial dose was 4.62 to 27.27 mg/kg/day and 4.29 to 30.00 mg/kg/day maintenance dose. The primary endpoint was reduction in seizure frequency from baseline at week 24. In the age group 1 month to 6 years (n=27) the number of seizures changed from 1 [range] [1-12] to 0 [0-2], in the age group 7 years to 12 years (n=77) the frequency changed from 1 [1-22] to 0 [0-1] and in the age group 13 to 19 years (n=152), the frequency changed from 1 [1-32] to 0 [0-3]. No specific safety concerns in the paediatric patients were identified. Data supporting benefit/risk from the study regarding children under the age of 6 are inconclusive.

Paediatric population

Based on the data from the randomised controlled trials, the use of oxcarbazepine is not recommended in children below the age of 6 since safety and efficacy have not been adequately demonstrated.

Two randomised, rater-blinded, dose-controlled efficacy studies (Study 2339 and Study 2340) were conducted in paediatric patients aged 1 month to <17 years of age (n=31 patients aged 6 to <17 years; n=189 patients aged <6 years old). In addition, a number of open-label studies that enrolled children were conducted. In general, the safety profile of oxcarbazepine in younger children (<6 years old) was similar to that in older children (≥ 6 years old). However, in some studies in younger children (<4 years old) and older children (≥ 4 years old), a ≥ 5 -fold difference in the proportion of patients with convulsions (7.9% vs. 1.0%, respectively) and status epilepticus (5% vs. 1%, respectively) was observed.

5.3 PHARMACOKINETIC PROPERTIES

Absorption

Following oral administration of oxcarbazepine, oxcarbazepine is completely absorbed and extensively metabolised to its pharmacologically active metabolite (MHD).

After a single dose administration of 600mg oxcarbazepine to healthy male volunteers under fasted conditions, the mean $C_{\max}$ value of MHD was 34 $\mu\text{mol/l}$, with a corresponding median $t_{\max}$ of 4.5 hours.

In a mass balance study in man, only 2 % of total radioactivity in plasma was due to unchanged oxcarbazepine, approximately 70 % was due to MHD, and the remainder attributable to minor secondary metabolites which were rapidly eliminated.

Food has no effect on the rate and extent of absorption of oxcarbazepine, therefore, oxcarbazepine can be taken with or without food.

Distribution

The apparent volume of distribution of MHD is 49 litres.

Approximately 40 % of MHD, is bound to serum proteins, predominantly to albumin. Binding was independent of serum concentration within the therapeutically relevant range. Oxcarbazepine and MHD do not bind to alpha-1-acid glycoprotein.

Oxcarbazepine and MHD cross the placenta. Neonatal and maternal plasma MHD concentrations were similar in one case.

Biotransformation

Oxcarbazepine is rapidly reduced by cytosolic enzymes in the liver to MHD, which is primarily responsible for the pharmacological effect of oxcarbazepine. MHD is metabolised further by conjugation with glucuronic acid. Minor amounts (4 % of the dose) are oxidised to the pharmacologically inactive metabolite (10, 11-dihydroxy derivative, DHD).

Elimination

Oxcarbazepine is cleared from the body mostly in the form of metabolites which are predominantly excreted by the kidneys. More than 95 % of the dose appears in the urine, with less than 1 % as unchanged oxcarbazepine. Faecal excretion accounts for less than 4 % of the administered dose. Approximately 80 % of the dose is excreted in the urine either as glucuronides of MHD (49 %) or as unchanged MHD (27 %), whereas the inactive DHD accounts for approximately 3 % and conjugates of oxcarbazepine account for 13 % of the dose.

Oxcarbazepine is rapidly eliminated from the plasma with apparent half-life values between 1.3 and 2.3 hours. In contrast, the apparent plasma half-life of MHD averaged 9.3 ± 1.8 h.

Linearity and dose proportionality

Steady-state plasma concentrations of MHD are reached within 2 - 3 days in patients when oxcarbazepine is given twice a day. At steady-state, the pharmacokinetics of MHD are linear and show dose proportionality across the dose range of 300 to 2400 mg/day.

Special populations

Patients with hepatic impairment

The pharmacokinetics and metabolism of oxcarbazepine and MHD were evaluated in healthy volunteers and hepatically-impaired subjects after a single 900 mg oral dose. Mild to moderate hepatic impairment did not affect the pharmacokinetics of oxcarbazepine and MHD. Oxcarbazepine has not been studied in patients with severe hepatic impairment.

Patients with renal impairment

There is a linear correlation between creatinine clearance and the renal clearance of MHD. When oxcarbazepine is administered as a single 300 mg dose, in renally impaired patients (creatinine clearance < 30 mL/min) the elimination half-life of MHD is prolonged by 60-90 % (16 to 19 hours) with a two fold increase in AUC compared to adults with normal renal function (10 hours).

Paediatric population

The pharmacokinetics of oxcarbazepine were evaluated in clinical trials in paediatric patients taking oxcarbazepine in the dose range 10-60 mg/kg/day. Weight-adjusted MHD clearance decreases as age and weight increases approaching that of adults. The mean weight-adjusted clearance in children 4 to 12 years of age is approximately 40% higher than that of adults. Therefore, MHD exposure in these children is expected to be about two-thirds that of adults

when treated with a similar weight-adjusted dose. As weight increases, for patients 13 years of age and above, the weight adjusted MHD clearance is expected to reach that of adults.

Pregnancy

Data from a limited number of women indicate that MHD plasma levels may gradually decrease throughout pregnancy.

Elderly

Following administration of single (300 mg) and multiple doses (600 mg/day) of oxcarbazepine in elderly volunteers (60 - 82 years of age), the maximum plasma concentrations and AUC values of MHD were 30 % - 60 % higher than in younger volunteers (18 - 32 years of age). Comparisons of creatinine clearances in younger and elderly volunteers indicate that the difference was due to age-related reductions in creatinine clearance. No special dose recommendations are necessary because therapeutic doses are individually adjusted.

Gender

No gender related pharmacokinetic differences have been observed in children, adults, or the elderly.

6. NONCLINICAL PROPERTIES

6.1 ANIMAL TOXICOLOGY OR PHARMACOLOGY

Preclinical data indicated no special hazard for humans based on safety pharmacology and genotoxicity studies with oxcarbazepine and the pharmacologically active metabolite, monohydroxy derivative (MHD).

Evidence of nephrotoxicity was noted in repeated dose toxicity rat studies but not in dog or mice studies.

Immunotoxicity

Immunostimulatory tests in mice showed that MHD (and to a lesser extent oxcarbazepine) can induce delayed hypersensitivity.

Mutagenicity

Oxcarbazepine increased mutation frequencies in one Ames test in vitro in the absence of metabolic activation in one of five bacterial strains. Oxcarbazepine and MHD produced increases in chromosomal aberrations and/or polyploidy in the Chinese hamster ovary assay in vitro in the absence of metabolic activation. MHD was negative in the Ames test, and no mutagenic or clastogenic activity was found with either oxcarbazepine or MHD in V79 Chinese hamster cells in vitro. Oxcarbazepine and MHD were both negative for clastogenic or aneugenic effects (micronucleus formation) in an in vivo rat bone marrow assay.

Reproductive toxicity

In rats, fertility in both sexes was unaffected by oxcarbazepine at oral doses up to 150 mg/kg/day, at which there is no safety margin. Disruption of estrous cyclicity and reduced numbers of corpora lutea, implantations and live embryos were observed in female animals for MHD at doses comparable to those in humans.

Standard reproductive toxicity studies in rodents and rabbits revealed effects such as increases in the incidence of embryo-foetal mortality and/or some delay in antenatal and/or postnatal

growth of the offspring at maternally toxic dose levels. There was an increase in rat foetal malformations in one of the eight embryo-foetal toxicity studies, which were conducted with either oxcarbazepine or MHD, at doses which also caused maternal toxicity.

Carcinogenicity

In the carcinogenicity studies, liver (rats and mice), testicular and female genital tract granular cell (rats) tumours were induced in treated animals. The occurrence of liver tumours was most likely a consequence of the induction of hepatic microsomal enzymes; an inductive effect which, although it cannot be excluded, is weak or absent in patients treated with oxcarbazepine. Testicular tumours may have been induced by elevated luteinising hormone concentrations.

Due to the absence of such an increase in humans, these tumours are considered to be of no clinical relevance. A dose-related increase in the incidence of granular cell tumours of the female genital tract (cervix and vagina) was noted in the rat carcinogenicity study with MHD. These effects occurred at exposure levels comparable with the anticipated clinical exposure. The mechanism for the development of these tumours has not been fully elucidated but could be related to increased estradiol levels specific to the rat. The clinical relevance of these tumours is unclear.

7. DESCRIPTION

Oxcarbazepine Tablets 150 mg:

OXMAZETOL is supplied for oral administration, as a film coated tablet of Oxcarbazepine. Each film coated tablet of OXMAZETOL contains Oxcarbazepine 150mg.

Oxcarbazepine Tablets 300mg:

OXMAZETOL is supplied for oral administration, as a film coated tablet of Oxcarbazepine. Each film coated tablet of OXMAZETOL contains Oxcarbazepine 300mg.

Oxcarbazepine Tablets 450 mg:

OXMAZETOL is supplied for oral administration, as a film coated tablet of Oxcarbazepine. Each film coated tablet of OXMAZETOL contains Oxcarbazepine 450mg.

Oxcarbazepine Tablets 600mg:

OXMAZETOL is supplied for oral administration, as a film coated tablet of Oxcarbazepine. Each film coated tablet of OXMAZETOL contains Oxcarbazepine 600mg.

8. PHARMACEUTICAL PARTICULARS

8.1 INCOMPATIBILITIES

Not applicable.

8.2 SHELF-LIFE

Refer Pack

8.3 PACKAGING INFORMATION

Refer Pack

8.4 STORAGE CONDITION AND HANDING INSTRUCTIONS

Refer Pack

Version 5.0; Dated 4-May-26

Replaces 4.0; Dated 11th Sep 2023

9. PATIENT COUNSELLING INFORMATION

Kindly refer to section 4.4

10. DETAILS OF MANUFACTURER

Refer Pack for manufacturer details.

Marketed By

Abbott Healthcare Pvt. Ltd.

Angel Space, Bldg. D-4, Gala No. 1 to 3 &

11 to 13 Ground Floor, 101 to 106 &

111 to 116 First Floor, 201 to 206 & 211 to 216

2nd Floor, Pimplas, Dist. Thane, Bhiwandi - 421 302,

Maharashtra, India.

11. DETAILS OF PERMISSION OR LICENCE NUMBER WITH DATE

Refer Pack for Permission/License details

12. DATE OF REVISION

Version 5.0; Dated 4-May-26

For Product Complaints/Adverse events or Queries please write to

webmasterindia@abbott.com